

IDH2 Rabbit pAb

货号: B12311

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IHC IF/ICC IP ChIP
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:20 - 1:50 IP: 1:50 - 1:200 ChIP: 1:20 - 1:100
理论分子量	45kDa/50kDa
实测分子量	43KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse heart,Mouse kidney,Rat kidney
细胞定位	Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 193-452 of human IDH2 (NP_002159.2).
序列	KMVFTPKDGSQVKEWEVYNFPAGGVGMGMYNTDEISGFAHSCFQYAIQKKWPLYMSTKNTILKAYDGRFKDIFQEIFDK HYKTDFDKNKIWIYEHRLIDDMVAQVLKSSGGFVWACKNYDGDVQSDILAQQFGSLGLMTSVLVCPDGTKIEAAHGT VTRHYREHQGRPTSTNPIASIFAWTRGLEHRGKLDGNQDLIRFAQMLEKVCVETVESGAMTKDLAGCIHGLSNVKLNEH FLNTTDFLDTIKSNLDRALGRQ

靶点信息

研究背景	Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the mitochondria. It plays a role in intermediary metabolism and energy production. This protein may tightly associate or interact with the pyruvate dehydrogenase complex. Alternative splicing results in multiple transcript variants.
基因ID	3418
基因名	IDH2
Swiss	P48735
别名	IDH2;D2HGA2;ICD-M;IDH;IDHM;IDP;IDPM;mNADP-IDH

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn