

OGFR Rabbit pAb

货号: B12316

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Homo sapiens |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 71kDa/73kDa |
| 实测分子量 | 80KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HepG2,A-549,Mouse lung,Mouse brain,Rat brain |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 358-482 of human OGF R (NP_031372.2). |
| 序列 | LERSQGDEAGGHGEDRPEPLSPKESKKRKLELSRREQPPTEPGPQSASEVEKIALNLEGICALSQGSLRTGTQEVEGGQDPG EAVQPCRQPLGARVADKVRKRRKVDEGAGDSAASGGAQTLALA |

靶点信息

| | |
|-------|--|
| 研究背景 | The protein encoded by this gene is a receptor for opioid growth factor (OGF), also known as [Met(5)]-enkephalin. OGF is a negative regulator of cell proliferation and tissue organization in a variety of processes. The encoded unbound receptor for OGF has been localized to the outer nuclear envelope, where it binds OGF and is translocated into the nucleus. The coding sequence of this gene contains a polymorphic region of 60 nt tandem imperfect repeat units. Several transcripts containing between zero and eight repeat units have been reported. |
| 基因ID | 11054 |
| 基因名 | OGFR |
| Swiss | Q9NZT2 |
| 别名 | OGFR |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn