

CSF2 Rabbit pAb

货号: **B12354**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	16kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,A-549,Mouse thymus
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human CSF2 (NP_000749.2) .
序列	MWLQSLLLLGTVACSISAPARSPSPSTQPWEHVNAIQEARRLLNLSRDTAAEMNETVEVISEMFDLQEPTCLQTRLELYKQ GLRGSLTKLKGPLTMMASH

靶点信息

研究背景	The protein encoded by this gene is a cytokine that controls the production, differentiation, and function of granulocytes and macrophages. The active form of the protein is found extracellularly as a homodimer . This gene has been localized to a cluster of related genes at chromosome region 5q31, which is known t o be associated with interstitial deletions in the 5q- syndrome and acute myelogenous leukemia. Other g enes in the cluster include those encoding interleukins 4, 5, and 13.
------	---

基因ID	1437
基因名	CSF2
Swiss	P04141
别名	CSF2;GMCSF

产品验证



Western blot analysis of CSF2 expressed in Jurkat, A-549, Mouse thymus using CSF2 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn