

# **HMOX2** Rabbit pAb

货号**: B12427** 

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	32kDa/36kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,HT-29,Mouse testis
细胞定位	Endoplasmic reticulum,Microsome
纯化	Affinity purification

# 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 400-500 of human HMOX2 (P30519).
序列	

# 靶点信息

研究背景	Heme oxygenase, an essential enzyme in heme catabolism, cleaves heme to form biliverdin, which is sub sequently converted to bilirubin by biliverdin reductase, and carbon monoxide, a putative neurotransmitt er. Heme oxygenase activity is induced by its substrate heme and by various nonheme substances. Hem e oxygenase occurs as 2 isozymes, an inducible heme oxygenase-1 and a constitutive heme oxygenase-2. HMOX1 and HMOX2 belong to the heme oxygenase family. Several alternatively spliced transcript varia nts encoding three different isoforms have been found for this gene.
基因 <b>ID</b>	3163

基因名	HMOX2
Swiss	P30519
别名	HMOX2;HO-2

### 产品验证



Western blot analysis of HMOX2 expressed in MCF7,HT-29,Mouse testis using HMOX2 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn