

# **BMPR1B** Rabbit pAb

货号: **B12475** 

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Sus scrofa
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	56kDa/60kDa
实测分子量	51KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,SH-SY5Y,293T,Mouse brain,Rat brain
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


# 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-126 of human BMPR1 B (NP_001194.1).
序列	MLLRSAGKLNVGTKKEDGESTAPTPRPKVLRCKCHHHCPEDSVNNICSTDGYCFTMIEEDDSGLPVVTSGCLGLEGSDFQ CRDTPIPHQRRSIECCTERNECNKDLHPTLPPLKNRDFVDGPIHHR

靶点信息

研究背景	This gene encodes a member of the bone morphogenetic protein (BMP) receptor family of transmembran e serine/threonine kinases. The ligands of this receptor are BMPs, which are members of the TGF-beta su perfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transd uce their signals through the formation of heteromeric complexes of 2 different types of serine (threonine ) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II receptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. Mutation s in this gene have been associated with primary pulmonary hypertension. Several transcript variants encoding two different isoforms have been found for this gene.
基因 <b>ID</b>	658
基因名	BMPR1B
Swiss	O00238
别名	ALK-6;ALK6;AMDD;BDA1D;BDA2;CDw293;BMPR1B

# 产品验证



Western blot analysis of BMPR1B expressed in Mouse brain,Rat brain,HeLa using BMPR1B Rabbit pAb a t 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per I ane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn