

VASN Rabbit pAb

货号: B12504

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	71kDa
实测分子量	71kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse testis,Mouse lung,Rat lung
细胞定位	cell surface,extracellular exosome,extracellular space,mitochondrion,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 420-575 of human VASN (NP_612449.2).
序列	CHLGTRHHLACLCPEGFTGLYCESQMGQGTRPSPTPVTPRPPRSLTLGIEPVSPVSLRVGLQRYLQGSSVQLRSLRLTYRNL SGPDKRLVTLRLPASLAEYTVTQLRPNATYSVCMPLGPGRVPEGEEACGEAHTPPAVHSNHAPVTQAREGNLP

靶点信息

研究背景	Vasorin is a Type I membrane protein, which is predominantly expressed in vascular smooth muscle cells in a developmentally regulated pattern. The expression level of Vasorin can be down regulated during vessel repair after arterial injury. Vasorin binds to transforming growth factor beta (TGF-β) and attenuates TGF-β signaling in vitro.
基因ID	114990

基因名	VASN
Swiss	Q6EMK4
别名	VASN;SLITL2;vasorin

产品验证



Western blot analysis of VASN expressed in Mouse kidney,Rat lung,Mouse testis using VASN Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn