

Acetyl-Histone H3-K4 Rabbit pAb

货号: B12508

产品信息

反应	Human,Mouse,Rat,Other (Wide Range)
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus IF: Mus musculus
应用	WB IF/ICC ChIP
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:100 ChIP: 1:50 - 1:200
理论分子量	15kDa
实测分子量	15kDa/17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,C6
细胞定位	Chromosome,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic acetylated peptide around K4 of human Histone H3 (NP_003484.1).
序列	ARTKQ

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.
基因ID	8290
基因名	HIST3H3
Swiss	Q16695
别名	H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn