

# COL5A1 Rabbit pAb

货号: **B12528** 

## 产品信息

| 反应    | Mouse   |
|-------|---|
| 宿主    | Rabbit  |
| 克隆性   | Polyclonal  |
| 预测反应  | WB: Homo sapiens, Mus musculus IHC: Rattus norvegicus   |
| 应用    | WB  |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000   |
| 理论分子量 | 183kDa  |
| 实测分子量 | 180kDa  |
| 形式    | Liquid  |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated  |
| 阳性对照  | Mouse lung  |
| 细胞定位  | Secreted,extracellular matrix,extracellular space   |
| 纯化    | Affinity purification   |


## 抗原信息

| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 20-120 of human COL5A1 (NP_0000 84.3). |  |
|------|---|--|
| 序列   | LPPLLLLLLWAPPPSRAAQPADLLKVLDFHNLPDGITKTTGFCATRRSSKGPDVAYRVTKDAQLSAPTKQLYPASAFPEDFS ILTTVKAKKGSQAFLVSIY    |  |

靶点信息

| 研究背景         | This gene encodes an alpha chain for one of the low abundance fibrillar collagens. Fibrillar collagen molec ules are trimers that can be composed of one or more types of alpha chains. Type V collagen is found in t issues containing type I collagen and appears to regulate the assembly of heterotypic fibers composed of both type I and type V collagen. This gene product is closely related to type XI collagen and it is possible t hat the collagen chains of types V and XI constitute a single collagen type with tissue-specific chain comb inations. The encoded procollagen protein occurs commonly as the heterotrimer pro-alpha1(V)-pro-alpha1(V)-pro-alpha1(V)-pro-alpha2(V). Mutations in this gene are associated with Ehlers-Danlos syndrome, types I and II. Alte rnative splicing of this gene results in multiple transcript variants. |
|--------------|---|
| 基因 <b>ID</b> | 1289  |
| 基因名          | COL5A1  |
| Swiss        | P20908  |
| 别名           | COL5A1;EDSC   |

## 产品验证



Western blot analysis of COL5A1 expressed in Mouse lung using COL5A1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn