

# MLKL Rabbit pAb

货号: **B12572** 

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens WB, IF: Rattus norvegicus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	53kDa/54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,NCI-H460,Mouse liver,Rat testis
细胞定位	Cell membrane,Cytoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-62 of mouse MLKL (Q9 D2Y4).
序列	MDKLGQIIKLGQLIYEQCEKMKYCRKQCQRLGNRVHGLLQPLQRLQAQGKKNLPDDITAALG

靶点信息

研究背景	This gene belongs to the protein kinase superfamily. The encoded protein contains a protein kinase-like d omain; however, is thought to be inactive because it lacks several residues required for activity. This prot ein plays a critical role in tumor necrosis factor (TNF)-induced necroptosis, a programmed cell death proc ess, via interaction with receptor-interacting protein 3 (RIP3), which is a key signaling molecule in necropt osis pathway. Inhibitor studies and knockdown of this gene inhibited TNF-induced necrosis. High levels of this protein and RIP3 are associated with inflammatory bowel disease in children. Alternatively spliced tra nscript variants have been described for this gene.
基因ID	74568
基因名	
Swiss	Q9D2Y4
别名	MLKL;hMLKL

## 产品验证



Western blot analysis of MLKL expressed in Mouse liver,Rat testis,Jurkat using MLKL Rabbit pAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn