

ZIP8 Rabbit pAb

货号: **B12659**

产品信息

| 反应 | Human,Mouse |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB, IP: Homo sapiens, Mus musculus WB: Bos taurus |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 43kDa/48kDa/49kDa |
| 实测分子量 | 50kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HL-60,NCI-H460,HeLa,Mouse liver |
| 细胞定位 | Membrane,Multi-pass membrane protein |
| 纯化 | Affinity purification |
| | |


抗原信息

| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 180-300 of human ZIP8 (NP_071437.3). |
|------|--|
| 序列 | QLIPEAFGFDPKVDSYVEKAVAVFGGFYLLFFFERMLKMLLKTYGQNGHTHFGNDNFGPQEKTHQPKALPAINGVTCYAN PAVTEANGHIHFDNVSVVSLQDGKKEPSSCTCLKGPKLSEI |

靶点信息

| This gene encodes a member of the SLC39 family of solute-carrier genes, which show structural character istics of zinc transporters. The encoded protein is glycosylated and found in the plasma membrane and m itochondria, and functions in the cellular import of zinc at the onset of inflammation. It is also thought to be the primary transporter of the toxic cation cadmium, which is found in cigarette smoke. Multiple transc ript variants encoding different isoforms have been found for this gene. Additional alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known. |
|---|
| 64116 |
| SLC39A8 |
| Q9C0K1 |
| SLC39A8;BIGM103;CDG2N;LZT-Hs6;PP3105;ZIP8 |
| |

产品验证



Western blot analysis of ZIP8 expressed in Mouse liver,HL-60,NCI-H460 using ZIP8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn