

# **MLKL Rabbit pAb**

货号: **B12713** 

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Rattus norvegicus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	53kDa/54kDa
实测分子量	54KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HT-29,Mouse liver
细胞定位	Cell membrane,Cytoplasm
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 369-471 of human MLKL (NP_689862.1).	
序列	DRVKSTAYLSPQELEDVFYQYDVKSEIYSFGIVLWEIATGDIPFQGCNSEKIRKLVAVKRQQEPLGEDCPSELREIIDECRAH DPSVRPSVDEILKKLSTFSK	

靶点信息

研究背景	This gene belongs to the protein kinase superfamily. The encoded protein contains a protein kinase-like d omain; however, is thought to be inactive because it lacks several residues required for activity. This prot ein plays a critical role in tumor necrosis factor (TNF)-induced necroptosis, a programmed cell death proc ess, via interaction with receptor-interacting protein 3 (RIP3), which is a key signaling molecule in necropt osis pathway. Inhibitor studies and knockdown of this gene inhibited TNF-induced necrosis. High levels of this protein and RIP3 are associated with inflammatory bowel disease in children. Alternatively spliced transcript variants have been described for this gene.
基因 <b>ID</b>	197259
基因名	MLKL
Swiss	Q8NB16
别名	hMLKL;MLKL

### 产品验证



Western blot analysis of MLKL expressed in HeLa,HT-29,Mouse liver using MLKL Rabbit pAb at 1:1000. S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blockin g buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn