

Ubiquitin Rabbit pAb

货号: **B12757**

产品信息


反应	Human,Mouse,Rat,Other (Wide Range)
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens IF: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	25kDa
实测分子量	20-150KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3,U-87MG
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-229 of human Ubiquiti n (NP_061828.1).
序列	MQIFVKTLTGKTITLEVEPSDTIENVKAKIQDKEGIPPDQQRLIFAGKQLEDGRTLSDYNIQKESTLHLVLRLRGGMQIFVKTL TGKTITLEVEPSDTIENVKAKIQDKEGIPPDQQRLIFAGKQLEDGRTLSDYNIQKESTLHLVLRLRGGMQIFVKTLTGKTITLE VEPSDTIENVKAKIQDKEGIPPDQQRLIFAGKQLEDGRTLSDYNIQKESTLHLVLRLRGGC

研究背景	This gene encodes ubiquitin, one of the most conserved proteins known. Ubiquitin has a major role in targ eting cellular proteins for degradation by the 26S proteosome. It is also involved in the maintenance of ch romatin structure, the regulation of gene expression, and the stress response. Ubiquitin is synthesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin moiety fused to an unrel ated protein. This gene consists of three direct repeats of the ubiquitin coding sequence with no spacer s equence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. An aberrant form of this protein has been detected in patients with Alzheimer's disease a nd Down syndrome. Pseudogenes of this gene are located on chromosomes 1, 2, 13, and 17. Alternative s plicing results in multiple transcript variants.
基因 ID	7314
基因名	UBB
Swiss	P0CG47
别名	UBB; HEL-S-50; ubiquitin B

产品验证



Western blot analysis of Ubiquitin expressed in NIH/3T3,U-87MG using Ubiquitin Rabbit pAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking b uffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn