

TIGAR Rabbit pAb

货号: **B12857**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	30kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse skeletal muscle,Rat skeletal muscle
细胞定位	cytoplasm,cytosol,mitochondrial outer membrane,nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human TIGAR (NP_06510 8.1).	
序列	LSELRAMAKAAREECPVFTPPGGETLDQVKMRGIDFFEFLCQLILKEADQKEQFSQGSPSNCLETSLAEIFPLGKNHSSKVN SDSGIPGLAASVLVVSHGA	

靶点信息

研究背景	This gene is regulated as part of the p53 tumor suppressor pathway and encodes a protein with sequence similarity to the bisphosphate domain of the glycolytic enzyme that degrades fructose-2,6-bisphosphate. The protein functions by blocking glycolysis and directing the pathway into the pentose phosphate shunt. Expression of this protein also protects cells from DNA damaging reactive oxygen species and provides s ome protection from DNA damage-induced apoptosis. The 12p13.32 region that includes this gene is par alogous to the 11q13.3 region.

基因 ID	57103
基因名	TIGAR
Swiss	Q9NQ88
别名	FR2BP; C12orf5

产品验证



Western blot analysis of TIGAR expressed in 293T,Mouse skeletal muscle,Rat skeletal muscle using TIGA R Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure tim e: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn