

PSPC1 Rabbit pAb

货号: B12919

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IP: Mouse ESC WB: Mus musculus
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:50 - 1:200
理论分子量	45kDa/58kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,A-549,U-87MG,Mouse brain
细胞定位	Cytoplasm,Nucleus,Nucleus matrix,Nucleus speckle,nucleolus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-70 of human PSPC1 (N P_001035879.1).
序列	MMLRGNLKQVRIEKNPARLRALESAVGESEPAAMALALAGEPAPPAPAPPEDHPDEEMGFTIDIKSF

靶点信息

研究背景	This gene encodes a nucleolar protein that localizes to punctate subnuclear structures that occur close to splicing speckles, known as paraspeckles. These paraspeckles are composed of RNA-protein structures that include a non-coding RNA, NEAT1/Men epsilon/beta, and the Drosophila Behavior Human Splicing family of proteins, which include the product of this gene and the P54NRB/NONO and PSF/SFPQ proteins. Paraspeckles may function in the control of gene expression via an RNA nuclear retention mechanism. The protein encoded by this gene is found in paraspeckles in transcriptionally active cells, but it localizes to unique cap structures at the nucleolar periphery when RNA polymerase II transcription is inhibited, or during telophase. Alternative splicing of this gene results in multiple transcript variants. A related pseudogene, which is also located on chromosome 13, has been identified.
基因ID	55269
基因名	PSPC1
Swiss	Q8WXF1
别名	PSPC1;PSP1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn