

Phospho-PDGFR beta-Y751 Rabbit pAb

货号: B12996

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	37kDa/123kDa
实测分子量	190kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Cell membrane,Cytoplasmic vesicle,Lysosome lumen,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息

抗原信息	A synthetic phosphorylated peptide around Y751 of human PDGFR beta (NP_002600.1).
序列	VDYVP


靶点信息

T.I	[477]	보보
117	「カルド	1 尽

This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth fact or family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth f actor bound to a receptor monomer determines whether the functional receptor is a homodimer or a hete rodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. This gen e is flanked on chromosome 5 by the genes for granulocyte-macrophage colony-stimulating factor and m acrophage-colony stimulating factor receptor; all three genes may be implicated in the 5-q syndrome. A t ranslocation between chromosomes 5 and 12, that fuses this gene to that of the translocation, ETV6, leuk emia gene, results in chronic myeloproliferative disorder with eosinophilia.

基因 ID	5159
基因名	PDGFRB
Swiss	P09619
别名	CD140B;IBGC4;IMF1;JTK12;KOGS;PDGFR;PDGFR-1;PDGFR1;PENTT;PDGF Receptor beta;PDGFRB;PDGFR be ta

产品验证



Western blot analysis of Phospho-PDGFR beta-Y751 expressed in NIH/3T3 using Phospho-PDGFR beta-Y751 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. E xposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn