

Phospho-p53-S46 Rabbit pAb

货号: B12998

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IP
推荐浓度	WB: 1:500 - 1:1000 IP: 1:50 - 1:100
理论分子量	23-43kDa
实测分子量	53KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Endoplasmic reticulum,Mitochondrion matrix,Nucleus,PML body
纯化	Affinity purification

抗原信息

抗原信息	A synthetic phosphorylated peptide around S46 of human p53 (NP_000537.3).
序列	MLSPD


靶点信息

砂	t	Ŧ	ť	7	ĭ	叧	ŧ

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and olig omerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metab olism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancer s such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937 277).

基因ID	7157
基因名	TP53
Swiss	P04637
别名	BCC7;LFS1;P53;TRP53;p53;TP53;BMFS5

产品验证



Western blot analysis of Phospho-p53-S46 expressed in HeLa using Phospho-p53-S46 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn