

Phospho-PDK1/PDPK1-S241 Rabbit pAb

货号: B12999

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	48kDa/50kDa/58kDa/60kDa/63kDa
实测分子量	60-70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Cell junction,Cell membrane,Cytoplasm,Nucleus,Peripheral membrane protein,focal adhesion
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S241 of human PDK1/PDPK1 (NP_002604.1).
序列	ANSFV

靶点信息

研究背景	Serine/threonine kinase which acts as a master kinase, phosphorylating and activating a subgroup of the AGC family of protein kinases. Plays a central role in the transduction of signals from insulin by providing the activating phosphorylation to PKB/AKT1, thus propagating the signal to downstream targets controlling cell proliferation and survival, as well as glucose and amino acid uptake and storage. Negatively regulates the TGF-beta-induced signaling . Activates PPARγ transcriptional activity and promotes adipocyte differentiation. Activates the NF-κB pathway via phosphorylation of IKKβ. The tyrosine phosphorylated form is crucial for the regulation of focal adhesions by angiotensin II. Controls proliferation, survival, and growth of developing pancreatic cells. Participates in the regulation of Ca(2+) entry and Ca(2+)-activated K(+) channels of mast cells. Essential for the motility of vascular endothelial cells (ECs) and is involved in the regulation of their chemotaxis. Plays a critical role in cardiac homeostasis by serving as a dual effector for cell survival and beta-adrenergic response. Plays an important role during thymocyte development by regulating the expression of key nutrient receptors on the surface of pre-T cells and mediating Notch-induced cell growth and proliferative responses. Provides negative feedback inhibition to toll-like receptor-mediated NF-κB activation in macrophages. Isoform 3 is catalytically inactive.
基因ID	5170
基因名	PDPK1
Swiss	O15530
别名	PDPK1;PDK1;PDPK2;PDPK2P;PRO0461

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn