

GFRA1 Rabbit pAb

货号: B13034

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: <i>Mus musculus</i>
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	50kDa/51kDa
实测分子量	51kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Mouse liver,Rat liver
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 25-280 of human GFRA1 (NP_665736.1).
序列	DRLDCVKASDQCLKEQSCSTKYRTLRQCVAGKETNFSLASGLEAKDECRSAMEALKQKSLYNCRCKRGMKKEKNCLRIY WSMYQSLQGNDLLEDSPYEPVNSRLSDIFRVVPFISVEHIPGNNCNDAAKACNLDDICKKYRSAYITPCFTSVSNDVCNR RKCHKALRQFFDKVPAKHSYGMFLCSRDIACTERRQTIVPVCSYEEREKPNCNLQDSCKTNYICRSRLADFFTNCQPES RSVSSCLKENYADC

靶点信息

研究背景	This gene encodes a member of the glial cell line-derived neurotrophic factor receptor (GDNFR) family of proteins. The encoded preproprotein is proteolytically processed to generate the mature receptor. Glial cell line-derived neurotrophic factor (GDNF) and neurturin (NTN) are two structurally related, potent neurotrophic factors that play key roles in the control of neuron survival and differentiation. This receptor is a glycosylphosphatidylinositol (GPI)-linked cell surface receptor for both GDNF and NTN, and mediates activation of the RET tyrosine kinase receptor. This gene is a candidate gene for Hirschsprung disease. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed.
基因ID	2674
基因名	GFRA1
Swiss	P56159
别名	GFRA1;GDNFR;GDNFRA;GFR-ALPHA-1;RET1L;RETL1;TRNR1

产品验证



Western blot analysis of GFRA1 expressed in MCF7, Mouse liver, Rat liver using GFRA1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn