

PTHLH Rabbit pAb

货号: **B13035**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	19kDa/20kDa/23kDa
实测分子量	26kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen
细胞定位	Cytoplasm,Nucleus,Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 74-130 of human PTHLH (NP_945316.1).
序列	ATSEVSPNSKPSPNTKNHPVRFGSDDEGRYLTQETNKVETYKEQPLKTPGKKKKGKP

靶点信息

研究背景	The protein encoded by this gene is a member of the parathyroid hormone family. This hormone, via its r eceptor, PTHR1, regulates endochondral bone development and epithelial-mesenchymal interactions duri ng the formation of the mammary glands and teeth. It is responsible for most cases of humoral hypercalc emia of malignancy, and mutations in this gene are associated with brachydactyly type E2 (BDE2). Altern atively spliced transcript variants have been found for this gene. There is also evidence for alternative tra nslation initiation from non-AUG (CUG and GUG) start sites, downstream of the initiator AUG codon, result ing in nuclear forms of this hormone.
------	--

基因ID	5744
基因名	PTHLH
Swiss	P12272
别名	PTHLH;BDE2;HHM;PLP;PTHR;PTHRP

产品验证



Western blot analysis of PTHLH expressed in Mouse spleen using PTHLH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn