

# Cathepsin L/V/K/H Rabbit pAb

货号: B13144

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Bos taurus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	37kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,293T
细胞定位	Lysosome
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 114-333 of human Cath epsin L/V/K/H (NP_001903.1).
序列	APRSVDWREKGYVTPVKNQGQCGSCWAFSATGALEGQMFRKTGRLISLSEQNLVDCSGPQGNEGCNGGLMDYAFQYV QDNGGLDSEESYPYEATEESCKYNPKYSVANDTGFVDIPKQEKALMKAVATVGPISVAIDAGHESFLFYKEGIYFEPDCSSE DMDHGVLVVGYGFESTESDNNKYWLVKNSWGEEWGMGGYVKMAKDRRNHCGIASAASYPTV

靶点信息

The protein encoded by this gene is a lysosomal cysteine proteinase that plays a major role in intracellula r protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a m ajor controlling element of neutrophil elastase activity. The encoded protein has been implicated in sever al pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria. This protein, which is a member of the peptidase C1 family, is a di mer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Multiple alternatively spliced transcript variants have been found for this gene.
1514
CTSL
P07711
CTSL;CATL;CTSL1;MEP

## 产品验证



Western blot analysis of Cathepsin L/V/K/H expressed in LO2,293T using Cathepsin L/V/K/H Rabbit pA b at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1 20s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn