

CPSF2 Rabbit pAb

货号: B13194

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:1000 - 1:2000
理论分子量	88kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,HeLa,U-937,Raji,LO2,Mouse kidney,Mouse liver
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 450-700 of human CPSF 2 (NP_059133.1).
序列	KGSFFKQAKKSYPMFPAPERIKWDEYGEIIPEDFLVPELQATEEEKSKLESGLTNGDEPMDQDLSDVPTKCISTTESIEIK ARVTYIDYEGRSDGDSIKKIINQMQRQLIIVHGPPEASQDLAECCEAFGGKDIKVYMPKLHETVDATSETHIYQVRLKDSL VSSLQFCKAKDAELAWIDGVLDMRVSKVDTGVILEEGELKDDGEDSEMVEAPSDSSVIAQQKAMKSLFGDDEKETGEES EIIPTL

靶点信息

研究背景	Component of the cleavage and polyadenylation specificity factor (CPSF complex that play a key role in p re-mRNA 3'-end formation, recognizing the AAUAAA signal sequence and interacting with poly(A polymer ase and other factors to bring about cleavage and poly(A addition. Involved in the histone 3' end pre-mR NA processing.
------	---

基因ID	53981
基因名	CPSF2
Swiss	Q9P2I0
别名	CPSF2;CPSF100

产品验证



Western blot analysis of CPSF2 expressed in Mouse kidney,U-251MG,HeLa using CPSF2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn