

DERL1 Rabbit pAb

货号: B13221

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Homo sapiens |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 26kDa/28kDa |
| 实测分子量 | 29kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | MCF7,Mouse liver |
| 细胞定位 | Endoplasmic reticulum membrane,Multi-pass membrane protein |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 182-251 of human DERL 1 (NP_077271.1). |
| 序列 | LYFFLMFRYPMDLGGRNFLSTPQFLYRWLPSSRRGGVSGFGVPPASMRRAADQNGGGGRHNWQGQFRLGDQ |

靶点信息

| | |
|------|---|
| 研究背景 | The protein encoded by this gene is a member of the derlin family. Members of this family participate in the ER-associated degradation response and retrotranslocate misfolded or unfolded proteins from the ER lumen to the cytosol for proteasomal degradation. This protein recognizes substrate in the ER and works in a complex to retrotranslocate it across the ER membrane into the cytosol. This protein may select cystic fibrosis transmembrane conductance regulator protein (CFTR) for degradation as well as unfolded proteins in Alzheimer's disease. Alternative splicing results in multiple transcript variants that encode different protein isoforms. |
|------|---|

| | |
|-------|---------------------------|
| 基因ID | 79139 |
| 基因名 | DERL1 |
| Swiss | Q9BUN8 |
| 别名 | DERL1;DER-1;DER1;derlin-1 |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn