

BRCC3 Rabbit pAb

货号: B13274

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	23kDa/28kDa/33kDa/36kDa
实测分子量	33kDa/36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,MCF7,Jurkat,293T,Mouse liver,Mouse kidney,Mouse brain,Rat liver
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-316 of human BRCC3 (NP_077308.1).
序列	MAVQVVQAVQAVHLESDAFLVCLNHALSTEKEEVMGLCIGELNDDTRSDSKFAYTGTEMRTVAEKVDAVRIVHIHSVILRR SDKRKDRVEISPEQLSAASTEERLAELTGRPMRVVGWYHSHPHITVWPSHVDVRTQAMYQMMDQGFVGLIFSCFIEDK NTKTGRVLYTCFQSIQAQKSSESLHGPRDFWSSSQHISIEGQKEERYERIEIPIHIVPHVTIGKVCLESAVELPKILCQEEQD AYRRIHSLTHLDSVTKIHNQSVFTKNLCSQMSAVSGPLLQWLEDRLEQNQQHLQELQKEELMQELSSLE

靶点信息

研究背景	This gene encodes a subunit of the BRCA1-BRCA2-containing complex (BRCC), which is an E3 ubiquitin ligase. This complex plays a role in the DNA damage response, where it is responsible for the stable accumulation of BRCA1 at DNA break sites. The component encoded by this gene can specifically cleave Lys 63-linked polyubiquitin chains, and it regulates the abundance of these polyubiquitin chains in chromatin. The loss of this gene results in abnormal angiogenesis and is associated with syndromic moyamoya, a cerebrovascular angiopathy. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 5.
基因ID	79184
基因名	BRCC3
Swiss	P46736
别名	BRCC3;BRCC36;C6.1A;CXorf53

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn