

# **OAZ1** Rabbit pAb

货号: **B13314** 

## 产品信息

| 反应    | Human, Mouse, Rat   |
|-------|---|
| 宿主    | Rabbit  |
| 克隆性   | Polyclonal  |
| 预测反应  | WB: Mus musculus  |
| 应用    | WB  |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000   |
| 理论分子量 | 25kDa   |
| 实测分子量 | 27kDa   |
| 形式    | Liquid  |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated  |
| 阳性对照  | MCF7,Mouse heart,Rat kidney   |
| 细胞定位  | cytoplasm,cytosol,nucleus   |
| 纯化    | Affinity purification   |


#### 抗原信息

| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-68 of human OAZ1 (N $P_004143.1$ ). |
|------|---|
| 序列   | MVKSSLQRILNSHCFAREKEGDKPSATIHASRTMPLLSLHSRGGSSSESSRVSLHCCSNPGPGPRWCS  |

靶点信息

| 研究背景         | The protein encoded by this gene belongs to the ornithine decarboxylase antizyme family, which plays a role in cell growth and proliferation by regulating intracellular polyamine levels. Expression of antizymes requires +1 ribosomal frameshifting, which is enhanced by high levels of polyamines. Antizymes in turn bind to and inhibit ornithine decarboxylase (ODC), the key enzyme in polyamine biosynthesis; thus, completing the auto-regulatory circuit. This gene encodes antizyme 1, the first member of the antizyme family, that has broad tissue distribution, and negatively regulates intracellular polyamine levels by binding to and targeting ODC for degradation, as well as inhibiting polyamine uptake. Antizyme 1 mRNA contains two potential in-frame AUGs; and studies in rat suggest that alternative use of the two translation initiation sites results in N-terminally distinct protein isoforms with different subcellular localization. Alternatively spliced transcript variants have also been noted for this gene. |
|--------------|---|
| 基因 <b>ID</b> | 4946  |
| 基因名          | OAZ1  |
| Swiss        | P54368  |
| 别名           | OAZ1;AZ1;AZI;OAZ  |

## 产品验证



Western blot analysis of OAZ1 expressed in MCF7,Mouse heart,Rat kidney using OAZ1 Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn