

TNFAIP6 Rabbit pAb

货号: B13438

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	31kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	extracellular region, extracellular space
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 110-277 of human TNFA IP6 (NP_009046.2).
序列	IIDYGIRLNRSERWDAYCYNPHAKECGGVFTDPKQIFKSPGFPNEYEDNQICYWHIRLKYGQRIHLSFLDFDLEDDPGCLAD YVEIYDSYDDVHGFVGRYCGDELPDDIISTGNVMTLKFLSDASVTAGGFQIKYVAMDPVSKSSQGKNTSTTSTGNKNFLA GRFSHL

靶点信息

研究背景	The protein encoded by this gene is a secretory protein that contains a hyaluronan-binding domain, and t hus is a member of the hyaluronan-binding protein family. The hyaluronan-binding domain is known to b e involved in extracellular matrix stability and cell migration. This protein has been shown to form a stabl e complex with inter-alpha-inhibitor (I alpha I), and thus enhance the serine protease inhibitory activity of I alpha I, which is important in the protease network associated with inflammation. This gene can be induced by proinflammatory cytokines such as tumor necrosis factor alpha and interleukin-1. Enhanced levels of this protein are found in the synovial fluid of patients with osteoarthritis and rheumatoid arthritis.
基因ID	7130
基因名	TNFAIP6
Swiss	P98066
别名	TNFAIP6;TSG-6;TSG6

产品验证



Western blot analysis of TNFAIP6 expressed in Mouse brain using TNFAIP6 Rabbit pAb at 1:1000. Secondar y antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn