

TADA3 Rabbit pAb

货号: B13460

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	41kDa/48kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,BxPC-3,HepG2,HeLa,Mouse testis,Mouse liver,Rat testis,Rat liver
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 270-369 of human TADA3 (NP_597814.1).
序列	EENIISPMEDSPIPDMSGKESGADGASTSPRNQNKPFVPHTKSLESRIKEELIAQGLLESEDRPAEDSEDEVLAELRKRQA ELKALSAHNRTKKHDLLR

靶点信息

研究背景	DNA-binding transcriptional activator proteins increase the rate of transcription by interacting with the transcriptional machinery bound to the basal promoter in conjunction with adaptor proteins, possibly by acetylation and destabilization of nucleosomes. The protein encoded by this gene is a transcriptional activator adaptor and a component of the histone acetyl transferase (HAT) coactivator complex which plays a crucial role in chromatin modulation and cell cycle progression. Along with the other components of the complex, this protein links transcriptional activators bound to specific promoters, to histone acetylation and the transcriptional machinery. The protein is also involved in the stabilization and activation of the p53 tumor suppressor protein that plays a role in the cellular response to DNA damage. Alternate splicing results in multiple transcript variants of this gene.
基因ID	10474
基因名	TADA3
Swiss	O75528
别名	TADA3;ADA3;NGG1;STAF54;TADA3L;hADA3

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn