

PER2 Rabbit pAb

货号: **B13629**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	45kDa/136kDa
实测分子量	136KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Nucleus,nucleolus,perinuclear region
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 390-565 of human PER2 (NP_073728.1).
序列	GQPFDYSPIRFRARNGEYITLDTSWSSFINPWSRKISFIIGRHKVRVGPLNEDVFAAHPCTEEKALHPSIQELTEQIHRLLLQP VPHSGSSGYGSLGSNGSHEHLMSQTSSSDSNGHEDSRRRRAEICKNGNKTKNRSHYSHESGEQKKKSVTEMQTNPPAE KKAVPAMEKDSLGV

靶点信息

研究背景 This gene is a member of the Period family of genes and is expressed in a circadian pattern hiasmatic nucleus, the primary circadian pacemaker in the mammalian brain. Genes in this components of the circadian rhythms of locomotor activity, metabolism, and behavior. This ulated by CLOCK/ARNTL heterodimers but then represses this upregulation in a feedback to CRY heterodimers to interact with CLOCK/ARNTL. Polymorphisms in this gene may increase ting certain cancers and have been linked to sleep disorders.	is family encode is gene is upreg loop using PER/
--	---

基因ID	8864
基因名	PER2
Swiss	O15055
别名	PER2;FASPS;FASPS1

产品验证



Western blot analysis of PER2 expressed in 293T using PER2 Rabbit pAb at 1:1000. Secondary antibody: H RP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn