

ZAP70 Rabbit pAb

货号: **B13706**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	35kDa/55kDa/69kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse thymus
细胞定位	Cell membrane,Cytoplasm,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human ZAP70 (NP_001070.2).
序列	MPDPA AHL PFFYGSISR AEEHLKLAGMADGLFLLRQCLRS LGGYVLSLVHDVRFHHFPIERQLNGTYAIAGGKAHCGPA ELCEFYSRDPDGLPCNLRKPCNRPSGLEPQPGVFDCLRDAMVRDYVRQ TWKLEGEALEQAIISQAPQVEKLIATTAHERMP WYHSSLTREEAERKLYSGAQT DGKFLLRPRKEQGTYALS LIYGKTVYHYLISQDKAGKYCIPEGTKFDTLWQLVEYLK LKAD GLIYCLKEACPSSASNASGAAAPTLPAHPSTLTHPQRRIDTLNSDGYTPEPARIT

靶点信息

研究背景	This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.
基因ID	7535
基因名	ZAP70
Swiss	P43403
别名	ZAP70;ADMIO2;IMD48;SRK;STCD;STD;TZK;ZAP-70

产品验证



Western blot analysis of ZAP70 expressed in Jurkat, Mouse thymus using ZAP70 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn