

CDK6 Rabbit pAb

货号: **B13725**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens, Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	36kDa
实测分子量	37kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,K-562
细胞定位	Cell projection,Cytoplasm,Nucleus,centrosome,cytoskeleton,microtubule organizing center,ruffle
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-326 of human CDK6 (NP_001250.1).
序列	MEKDGLCRADQQYECVAEIGEGAYGKVFKARDLKNGGRFVALKRVRVQTGEEGMPLSTIREVAVLRHLETFEHPNVVRLF DVCTVSRTDRETKLTLVFEHVDQDLTTYLDKVPEPGVPTETIKDMMFQLLRGLDFLHSHRVVHRDLKPQNILVTSSGQIKL ADFGLARIYSFQMALTSVVVTLWYRAPEVLLQSSYATPVDLWSVGCIFAEMFRRKPLFRGSSDVDQLGKILDVIGLPGEED WPRDVALPRQAFHSKSAQPIEKFVTDIDELGKDLLLKCLTFNPAKRISAYSALSHPYFQDLERCKENLDSHLPPSQNTSELN TA

研究背景	The protein encoded by this gene is a member of the cyclin-dependent protein kinase (CDK) family. CDK f amily members are highly similar to the gene products of Saccharomyces cerevisiae cdc28, and Schizosa ccharomyces pombe cdc2, and are known to be important regulators of cell cycle progression. This kinas e is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression and G1/S transition. The activity of this kinase first appears in mid-G1 phase, which is controlled by the re gulatory subunits including D-type cyclins and members of INK4 family of CDK inhibitors. This kinase, as well as CDK4, has been shown to phosphorylate, and thus regulate the activity of, tumor suppressor prot ein Rb. Expression of this gene is up-regulated in some types of cancer. Multiple alternatively spliced vari ants, encoding the same protein, have been identified.
基因ID	1021
基因名	CDK6
Swiss	Q00534
别名	CDK6;MCPH12;PLSTIRE

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn