

# **SMARCC2** Rabbit pAb

货号**: B13771** 

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	124kDa/126kDa/132kDa
实测分子量	171kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Mouse spinal cord
细胞定位	Nucleus
纯化	Affinity purification

#### 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human SMARCC2 (NP_62 0706.1).
序列	PKLLGKLKDIIKRHQGTVTEDKNNASHVVYPVPGNLEEEEWVRPVMKRDKQVLLHWGYYPDSYDTWIPASEIEASVEDAP TPEKPRKVHAKWILDTDTFNE

### 靶点信息

研究背景	The protein encoded by this gene is a member of the SWI/SNF family of proteins, whose members display helicase and ATPase activities and which are thought to regulate transcription of certain genes by alterin g the chromatin structure around those genes. The encoded protein is part of the large ATP-dependent c hromatin remodeling complex SNF/SWI and contains a predicted leucine zipper motif typical of many tran scription factors. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	6601
基因名	SMARCC2
Swiss	Q8TAQ2
别名	SMARCC2;BAF170;CRACC2;Rsc8

# 产品验证



Western blot analysis of SMARCC2 expressed in U-87MG, Mouse spinal cord using SMARCC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn