

# LAMC1 Rabbit pAb

货号: **B13830** 

### 产品信息


反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200
理论分子量	177kDa
实测分子量	220KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,Mouse heart,Mouse kidney,Rat heart
细胞定位	basement membrane,endoplasmic reticulum lumen,extracellular exosome,extracellular region,extracellul ar space
纯化	Affinity purification

# 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1300-1609 of human LA MC1 (NP_002284.3).
序列	IDQKLKDYEDLREDMRGKELEVKNLLEKGKTEQQTADQLLARADAAKALAEEAAKKGRDTLQEANDILNNLKDFDRRVND NKTAAEEALRKIPAINQTITEANEKTREAQQALGSAAADATEAKNKAHEAERIASAVQKNATSTKAEAERTFAEVTDLDNEV NNMLKQLQEAEKELKRKQDDADQDMMMAGMASQAAQEAEINARKAKNSVTSLLSIINDLLEQLGQLDTVDLNKLNEIEG TLNKAKDEMKVSDLDRKVSDLENEAKKQEAAIMDYNRDIEEIMKDIRNLEDIRKTLPSGCFNTPSIEKP

研究背景	Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of base ment membranes. They have been implicated in a wide variety of biological processes including cell adhe sion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins, composed of 3 no n identical chains: laminin alpha, beta and gamma. Some of the chains have been shown to differ with re spect to their tissue distribution, presumably reflecting diverse functions in vivo. This gene encodes the g amma chain isoform laminin, gamma 1. The gamma 1 chain, formerly thought to be a beta chain, contain s structural domains similar to beta chains, however, lacks the short alpha region separating domains I a nd II. The structural organization of this gene also suggested that it had diverged considerably from the b eta chain genes. Embryos of transgenic mice in which both alleles of the gamma 1 chain gene were inacti vated by homologous recombination, lacked basement membranes, indicating that laminin, gamma 1 ch ain is necessary for laminin heterotrimer assembly. It has been inferred by analogy with the strikingly sim ilar 3' UTR sequence in mouse laminin gamma 1 cDNA, that multiple polyadenylation sites are utilized in human to generate the 2 different sized mRNAs (5.5 and 7.5 kb) seen on Northern analysis.
基因ID	3915
基因名	LAMC1
Swiss	P11047
别名	LAMC1;LAMB2

## 产品验证



Western blot analysis of LAMC1 expressed in Mouse heart,Rat heart,LO2 using LAMC1 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn