

# **ORC2** Rabbit pAb

## 货号**: B13863**

### 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens IP: Homo sapiens
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	65kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T
细胞定位	Nucleus
纯化	Affinity purification


#### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 458-577 of human ORC 2 (NP_006181.1).	
序列	TSYENSLLVKQSGSLPLSSLTHVLRSLTPNARGIFRLLIKYQLDNQDNPSYIGLSFQDFYQQCREAFLVNSDLTLRAQLTEFR DHKLIRTKKGTDGVEYLLIPVDNGTLTDFLEKEEEEA	

#### 靶点信息

研究背景	The origin recognition complex (ORC) is a highly conserved six subunits protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast demonstrated that ORC binds specific ally to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. The protein encoded by this gene is a subunit of the ORC complex. This protein n forms a core complex with ORC3, -4, and -5. It also interacts with CDC45 and MCM10, which are protein s known to be important for the initiation of DNA replication. This protein has been demonstrated to specifically associate with the origin of replication of Epstein-Barr virus in human cells, and is thought to be re quired for DNA replication from viral origin of replication. Alternatively spliced transcript variants have be en found, one of which is a nonsense-mediated mRNA decay candidate.
基因ID	4999
基因名	ORC2
Swiss	Q13416
别名	ORC2;ORC2L

#### 产品验证



Western blot analysis of ORC2 expressed in HeLa,293T using ORC2 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn