

# **ABCB7 Rabbit pAb**

货号: **B13947** 

### 产品信息


反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	78kDa/82kDa
实测分子量	83kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,293T,Mouse heart
细胞定位	Mitochondrion inner membrane, Multi-pass membrane protein
纯化	Affinity purification

### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 503-753 of human ABCB7 (NP_004290.2).
序列	AIVGGSGSGKSTIVRLLFRFYEPQKGSIYLAGQNIQDVSLESLRRAVGVVPQDAVLFHNTIYYNLLYGNISASPEEVYAVAKL AGLHDAILRMPHGYDTQVGERGLKLSGGEKQRVAIARAILKDPPVILYDEATSSLDSITEETILGAMKDVVKHRTSIFIAHRLS TVVDADEIIVLDQGKVAERGTHHGLLANPHSIYSEMWHTQSSRVQNHDNPKWEAKKENISKEEERKKLQEEIVNSVKGCG NCSC

研究背景	The membrane-associated protein encoded by this gene is a member of the superfamily of ATP-binding c assette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular me mbranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN2 0, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance as well as antigen presentation. This gene encodes a half-transporter involved in the transport of heme from the mitochondria to the cytosol. With iron/sulfur cluster precursors as its substrates, this protein may play a role in metal homeostasis. Mutations in this gene have been associate d with mitochondrial iron accumulation and isodicentric (X)(q13) and sideroblastic anemia. Alternatively s pliced transcript variants encoding multiple isoforms have been observed for this gene.
基因 <b>ID</b>	22
基因名	ABCB7
Swiss	075027
别名	ABCB7;ABC7;ASAT;Atm1p;EST140535

### 产品验证



Western blot analysis of ABCB7 expressed in A-431,293T,Mouse heart using ABCB7 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn