

TAPBP Rabbit pAb

货号: B14036

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Oryctolagus cuniculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	38kDa/43kDa/47kDa/53kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,Raji,M21,Mouse liver
细胞定位	Endoplasmic reticulum membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-260 of human TAPBP (NP_757345.2).
序列	VEDASGKGLAKRPGALLLRQGPGEPPRPDLDPELYLSVHDPAGALQAAFRRYPRGAPAPHCEMSRLFVPLPASAKWASGL TPAQNCPRALDGAWLMVSISSPVLSLSSLLRPQPEPQQEPVLITMATVVLTVLTHTPAPRVRLGQDALLDLSFAYMPPTSEA ASSLAPGPPPFGLEWRRQHLGKGHLLLAATPGLNGQMPAAQEGAVAFAAWDDDEPWGPWTGNGTFWLPT

靶点信息

研究背景	This gene encodes a transmembrane glycoprotein which mediates interaction between newly assembled major histocompatibility complex (MHC) class I molecules and the transporter associated with antigen processing (TAP), which is required for the transport of antigenic peptides across the endoplasmic reticulum membrane. This interaction is essential for optimal peptide loading on the MHC class I molecule. Up to four complexes of MHC class I and this protein may be bound to a single TAP molecule. This protein contains a C-terminal double-lysine motif (KKKAE) known to maintain membrane proteins in the endoplasmic reticulum. This gene lies within the major histocompatibility complex on chromosome 6. Alternative splicing results in three transcript variants encoding different isoforms.
基因ID	6892
基因名	TAPBP
Swiss	O15533
别名	TAPBP;NGS17;TAPA;TPN;TPSN;tapasin

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn