

# ZP3 Rabbit pAb

# 货号**: B14057**

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	41kDa/47kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cell membrane, Secreted, Single-pass type I membrane protein, extracellular matrix, extracellular space
纯化	Affinity purification

#### 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human ZP3 (NP_0011038 24.1).	
序列	AFKVPRPGPDTLQFTVDVFHFANDSRNMIYITCHLKVTLAEQDPDELNKACSFSKPSNSWFPVEGSADICQCCNKGDCGT PSHSRRQPHVMSQWSRSASRN	

#### 靶点信息

研究背景	The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is composed primarily of three or four glycoproteins with various functions during fertilization and preimplantation dev elopment. The protein encoded by this gene is a structural component of the zona pellucida and functions in primary binding and induction of the sperm acrosome reaction. The nascent protein contains a N-termi nal signal peptide sequence, a conserved ZP domain, a C-terminal consensus furin cleavage site, and a tr ansmembrane domain. It is hypothesized that furin cleavage results in release of the mature protein from the plasma membrane for subsequent incorporation into the zona pellucida matrix. However, the require ment for furin cleavage in this process remains controversial based on mouse studies. A variation in the la st exon of this gene has previously served as the basis for an additional ZP3 locus; however, sequence an d literature review reveals that there is only one full-length ZP3 locus in the human genome. Another locu s encoding a bipartite transcript designated POMZP3 contains a duplication of the last four exons of ZP3, i ncluding the above described variation, and maps closely to this gene.
基因ID	7784
基因名	ZP3
Swiss	P21754
别名	ZP3;ZP3A;ZP3B;ZPC;Zp-3

## 产品验证



Western blot analysis of ZP3 expressed in HeLa using ZP3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry mi lk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn