

EED Rabbit pAb

货号: B14084

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100
理论分子量	45kDa/50kDa/53kDa
实测分子量	50-75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,MCF7,A-549,K-562,Mouse liver,Mouse lung,Rat lung
细胞定位	Chromosome,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human EED (NP_003788.2).
序列	MSEREVSTAPAGTDMPAAKKQKLSSDENSENPDLSGDENDDAVSIESGTNTERPDTPTNTPNAPGRKSWGKGKWKSKKC KYSFKCVNSLKEDHNQPLFGVQ

靶点信息

研究背景	This gene encodes a member of the Polycomb-group (PcG) family. PcG family members form multimeric p rotein complexes, which are involved in maintaining the transcriptional repressive state of genes over suc cessive cell generations. This protein interacts with enhancer of zeste 2, the cytoplasmic tail of integrin b eta7, immunodeficiency virus type 1 (HIV-1) MA protein, and histone deacetylase proteins. This protein m ediates repression of gene activity through histone deacetylation, and may act as a specific regulator of i ntegrin function. Two transcript variants encoding distinct isoforms have been identified for this gene.
------	---

基因ID	8726
基因名	EED
Swiss	O75530
别名	EED;HEED;WAIT1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn