

# **BRCA1** Rabbit pAb

货号: **B14190** 

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	7kDa/78-85kDa/202-210kDa
实测分子量	7kDa/78-85kDa/202-210kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Chromosome,Cytoplasm,Cytoplasm,Nucleus
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 400-699 of human BRCA1 (NP_009230.2).
序列	LDDRWYMHSCSGSLQNRNYPSQEELIKVVDVEEQQLEESGPHDLTETSYLPRQDLEGTPYLESGISLFSDDPESDPSEDR APESARVGNIPSSTSALKVPQLKVAESAQSPAAAHTTDTAGYNAMEESVSREKPELTASTERVNKRMSMVVSGLTPEEFML VYKFARKHHITLTNLITEETTHVVMKTDAEFVCERTLKYFLGIAGGKWVVSYFWVTQSIKERKMLNEHDFEVRGDVVNGRN HQGPKRARESQDRKIFRGLEICCYGPFTNMPTGCPPNCGCAARCLDRGQWLPCNWADV

靶点信息

研究背景	This gene encodes a nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The encoded protein combines with other tumor suppressors, DNA damage s ensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associ ated genome surveillance complex (BASC). This gene product associates with RNA polymerase II, and thr ough the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. Mutations in this gene are responsible for approximately 40% of inherited breast cancers and more than 80% of inherited breast an d ovarian cancers. Alternative splicing plays a role in modulating the subcellular localization and physiolo gical function of this gene. Many alternatively spliced transcript variants, some of which are disease-assoc iated mutations, have been described for this gene, but the full-length natures of only some of these vari ants has been described. A related pseudogene, which is also located on chromosome 17, has been identified.
基因 <b>ID</b>	672
基因名	BRCA1
Swiss	P38398
别名	BRCA1;BRCA1;BRCC1;BROVCA1;FANCS;IRIS;PNCA4;PPP1R53;PSCP;RNF53

### 产品验证



Western blot analysis of BRCA1 expressed in human esophagus cancer, mouse lung, rat liver using BRCA1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn