

MCM2 Rabbit pAb

货号: B14209

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	101kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,SH-SY5Y,Raji,A-549,HeLa,HL-60,Mouse thymus
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 360-700 of human MCM 2 (NP_004517.2).
序列	PFEVNMEETIQNYQRIRIQESPGKVAAGRLPRSKDAILLADLVDCKPGDEIELTGIYHNNDGSLNTANGFPVFATVILAN HVAKKDNKVAVGELTDEDVKMITSLSKDQQIGEKIFASIAPSIYGHEDIKRLALALFGGEPKNPGGKHKVRGDINVLLCGD PGTAKSQFLKYIEKVSSRAIFTTGQASAVGLTAYVQRHPVSREWTLAAGALVLADRGVCLIDEFDKMNDQDRTSIHEAM EQQSISISKAGIVTSLQARCTVIAAANPIGGRYDPSLTFSENVDLTEPIISRFDILCVVRDTPVQDEMLARFVVGSHVRHH PSNKEEEGLANGS

靶点信息

研究背景	The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are involved in the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. This protein forms a complex with MCM4, 6, and 7, and has been shown to regulate the helicase activity of the complex. This protein is phosphorylated, and thus regulated by, protein kinases CDC2 and CDC7. Multiple alternatively spliced transcript variants have been found, but the full-length nature of some variants has not been defined.
基因ID	4171
基因名	MCM2
Swiss	P49736
别名	MCM2;BM28;CCNL1;CDCL1;D3S3194;DFNA70;MITOTIN;cdc19

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn