

PI3 Kinase p85 alpha Rabbit pAb

货号: B14312

产品信息

| | |
|-------|--|
| 反应 | Human,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Homo sapiens |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:1000 - 1:5000 IHC: 1:50 - 1:200 |
| 理论分子量 | 42kDa/49kDa/53kDa/83kDa/84kDa |
| 实测分子量 | 85KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-87MG |
| 细胞定位 | cell-cell junction,cis-Golgi network,cytoplasm,cytosol,nucleus,perinuclear region of cytoplasm,plasma membrane |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 125-320 of human PI3 Kinase p85 alpha (NP_852664.1). |
| 序列 | PDIAPLLIKLVEAIEKKGLECSTLYRTQSSSNLAELRQLLDCDTPSVDLEMIDVHVLADAFKRYLLDLPNPVIPAAVYSEMISL APEVQSSEYIQLLKKLIRSPSIPHQYWLTQYLLKHFFKLSQTSSKNLLNARVLSEIFSPMLFRFSAASSDNTENLIKVIEILIST EWNERQPAPALPPKPKPTTVANN |

靶点信息

| | |
|-------|---|
| 研究背景 | Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit. Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in four transcript variants encoding different isoforms. |
| 基因ID | 5295 |
| 基因名 | PIK3R1 |
| Swiss | P27986 |
| 别名 | AGM7;GRB1;IMD36;p85;p85-ALPHA;PI3 Kinase p85 alpha;PIK3R1 |

产品验证



Western blot analysis of PI3 Kinase p85 alpha expressed in U-87MG using PI3 Kinase p85 alpha Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s

实验步骤

访问官网浏览详情: www.ablybio.cn