

VHL Rabbit pAb

货号: **B14335**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	18kDa/19kDa/24kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat brain
细胞定位	Cytoplasm,Membrane,Nucleus,Nucleus,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human VHL (NP_937799.1).	
序列	MPRRAENWDEAEVGAEEAGVEEYGPEEDGGEESGAEESGPEESGPEELGAEEEMEAGRPRPVLRSVNSREPSQVIFCNR SPRVVLPVWLNFDGEPQPYPT	

靶点信息

研究背景	Von Hippel-Lindau syndrome (VHL) is a dominantly inherited familial cancer syndrome predisposing to a variety of malignant and benign tumors. A germline mutation of this gene is the basis of familial inheritan ce of VHL syndrome. The protein encoded by this gene is a component of the protein complex that includ es elongin B, elongin C, and cullin-2, and possesses ubiquitin ligase E3 activity. This protein is involved in the ubiquitination and degradation of hypoxia-inducible-factor (HIF), which is a transcription factor that pl ays a central role in the regulation of gene expression by oxygen. RNA polymerase II subunit POLR2G/RPB 7 is also reported to be a target of this protein. Alternatively spliced transcript variants encoding distinct i soforms have been observed.
基因 ID	7428
基因名	VHL
Swiss	P40337
别名	VHL;HRCA1;RCA1;VHL1;pVHL;PVHL

产品验证



Western blot analysis of VHL expressed in Mouse brain,Rat brain using VHL Rabbit pAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn