

VEGFA Rabbit pAb

货号**: B14373**

产品信息

| 反应 | Human,Mouse,Rat |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Homo sapiens |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 15-27kDa/34-45kDa |
| 实测分子量 | 16KDa/34KDa/45KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | A-549,Mouse liver,Rat lung |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |

抗原信息

| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 100-191 of human VEGFA (NP_0011 65097.1). | |
|------|--|--|
| 序列 | SNITMQIMRIKPHQGQHIGEMSFLQHNKCECRPKKDRARQENPCGPCSERRKHLFVQDPQTCKCSCKNTDSRCKARQLEL NERTCRCDKPRR | |

靶点信息

| 研究背景 | This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, whic h exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascul ar endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of thi s gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this p rotein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atheroscl erosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in addit ional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alter native in-frame translation termination codon via a stop codon readthrough mechanism, and that this isof orm is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a s mall upstream open reading frame, which is located within an internal ribosome entry site. |
|-------|---|
| 基因ID | 7422 |
| 基因名 | VEGFA |
| Swiss | P15692 |
| 别名 | VEGFA;MVCD1;VEGF;VPF;L VEGFA;VEGF A |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn