

# CYP7B1 Rabbit pAb

货号: B14407

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	58kDa
实测分子量	55KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431
细胞定位	Endoplasmic reticulum membrane,Microsome membrane,Peripheral membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human CYP7B1 (NP_0048 11.1).	
序列	VIVCDNNKFISELRDDFLKFDDKFAYLVSNIPIELLGNVKSIREKIIKCFSSEKLAKMQGWSEVFQSRQDVLEKYYVHEDLEIG AHHLGFLWASVANTIPT	

#### 靶点信息

研究背景	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 pro teins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of c holesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the first rea ction in the cholesterol catabolic pathway of extrahepatic tissues, which converts cholesterol to bile acids. This enzyme likely plays a minor role in total bile acid synthesis, but may also be involved in the develop ment of atherosclerosis, neurosteroid metabolism and sex hormone synthesis.

基因ID	9420
基因名	CYP7B1
Swiss	O75881
别名	CYP7B1;CBAS3;CP7B;SPG5A

## 产品验证



Western blot analysis of CYP7B1 expressed in A-431 using CYP7B1 Rabbit pAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn