

FOXO1/FOXO3A/FOXO4 Rabbit pAb

货号: B14446

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa,Mouse liver,Mouse skeletal muscle,Mouse lung,Rat liver
细胞定位	cytoplasm,cytosol,mitochondrion,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide of human FOXO1/FOXO3A/FOXO4 (NP_002006.2).
序列	

靶点信息

研究背景	The Forkhead family of transcription factors mediates many aspects of physiology, including stress response, metabolism, commitment to apoptosis, and development. The Forkhead box subfamily O (FoxO) proteins have garnered particular interest due to their involvement in the modulation of cancer, metabolism and cardiovascular biology. In mammals, four subfamily members have been identified: FOXO1 (previously also known as FKHR), FOXO3 (also known as FKHRL1), FOXO4 (also known as AFX), and FOXO6.
基因ID	2308,2309,4303

基因名	FOXO1,FOXO3,FOXO4
Swiss	Q12778,O43524,P98177
别名	

产品验证



Western blot analysis of FOXO1/FOXO3A/FOXO4 expressed in Mouse liver,Rat liver,293T using FO XO1/FOXO3A/FOXO4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detectio n: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn