

NCAM1 / CD56 Rabbit pAb

货号: B14879

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mouse lung tumor, Mouse C3 primary cells IHC: Mouse normal liver, Mouse lung tumor, Mouse C3 primary cells
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:100
理论分子量	40kDa/73kDa/80kDa/83kDa/93kDa/94kDa
实测分子量	120kDa/140kDa/180kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain tissue
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 759-858 of human NCA M1 / CD56 (NP_851996.2).
序列	LCGKAGPGAKGKDMEEGKAAFSKDESKEPIVEVRTEEERTPNHDGGKHTEPNETTPLTEPEKGPVEAKPECQETETKPAPA EVKTVPNDATQTKENESKA

靶点信息

研究背景	This gene encodes a cell adhesion protein which is a member of the immunoglobulin superfamily. The en coded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein has been shown to be involved in development of the nervous sy stem, and for cells involved in the expansion of T cells and dendritic cells which play an important role in i mmune surveillance. Alternative splicing results in multiple transcript variants.
基因 ID	4684
基因名	NCAM1
Swiss	P13591
别名	CD56;MSK39;NCAM1;NCAM1;CD56

产品验证



Western blot analysis of NCAM1 / CD56 expressed in Mouse brain tissue using NCAM1 / CD56 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lan e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn