

Phospho-MDM2-S166 Rabbit pAb

货号: B14883

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Adenocarcinoma human alveolar basal epithelial cells IHC: Adenocarcinoma human alveolar basal epithelial cells
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	11-14kDa/24-55kDa
实测分子量	95kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293
细胞定位	Cytoplasm,Nucleus,nucleolus,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around S166 of human MDM2 (NP_002383.2).
序列	TSSRR

靶点信息

研究背景	This gene encodes a nuclear-localized E3 ubiquitin ligase. The encoded protein can promote tumor formation by targeting tumor suppressor proteins, such as p53, for proteasomal degradation. This gene is itself transcriptionally-regulated by p53. Overexpression or amplification of this locus is detected in a variety of different cancers. There is a pseudogene for this gene on chromosome 2. Alternative splicing results in a multitude of transcript variants, many of which may be expressed only in tumor cells.
基因ID	4193

基因名	MDM2
Swiss	Q00987
别名	MDM2;ACTFS;HDMX;hdm2

产品验证



Western blot analysis of Phospho-MDM2-S166 expressed in 293 using Phospho-MDM2-S166 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn