

MTTP Rabbit pAb

货号: B14890

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: rat liver , Mouse liver
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	16kDa/99kDa
实测分子量	99kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Rat small intestine
细胞定位	Endoplasmic reticulum
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-130 of human MTTP (NP_000244.2).
序列	MILLAVLFLCFISSYSASVKGHTTGLSLNNDRLYKLTYSTEVLLDRGKGKLQDSVGYRISSNVDVALLWRNPDGDDDQLIQI TMKDVNVENVNQQRGEKSIFKGKSPSKIMGKENLEALQRPTLLHLIHG

靶点信息

研究背景	MTTP encodes the large subunit of the heterodimeric microsomal triglyceride transfer protein. Protein disul fide isomerase (PDI) completes the heterodimeric microsomal triglyceride transfer protein, which has bee n shown to play a central role in lipoprotein assembly. Mutations in MTP can cause abetalipoproteinemia.
基因ID	4547

基因名	MTTP
Swiss	P55157
别名	MTTP;ABL;MTP

产品验证



Western blot analysis of MTTP expressed in Mouse liver,Rat small intestine using MTTP Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bloc king buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn