

HTRA3 Rabbit pAb

货号: **B14891**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: human cells IHC: Rat bone tissue
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	37kDa/48kDa
实测分子量	49KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,SKOV3,Mouse brain
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 334-453 of human HTRA3 (NP_444272.1).
序列	SDRITRFLTEFQDKQIKDWKKRFIGIRMRTITPSLVDELKASNPDFPEVSSGIYVQEVAPNSPSQRGGIQDGDIIVKVNGRPL VDSSELQEAVLTESPLLLEVRRGNDDLLFSIAPEVVM

靶点信息

研究背景	Serine protease that cleaves beta-casein/CSN2 as well as several extracellular matrix (ECM proteoglycans such as decorin/DCN, biglycan/BGN and fibronectin/FN1. Inhibits signaling mediated by TGF-beta family p roteins possibly indirectly by degradation of these ECM proteoglycans (By similarity. May act as a tumor s uppressor. Negatively regulates, in vitro, trophoblast invasion during placental development and may be involved in the development of the placenta in vivo. May also have a role in ovarian development, granul osa cell differentiation and luteinization.
基因 ID	94031
基因名	HTRA3
Swiss	P83110
别名	HTRA3;Prsp;Tasp

产品验证



Western blot analysis of HTRA3 expressed in 293T,SKOV3,Mouse brain using HTRA3 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn