

# **PROCR Rabbit pAb**

货号**: B14903** 

## 产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: HepG2
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	26kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,HepG2,HeLa,293T,Mouse kidney
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 18-210 of human PROC R (NP_006395.2).
序列	SQDASDGLQRLHMLQISYFRDPYHVWYQGNASLGGHLTHVLEGPDTNTTIIQLQPLQEPESWARTQSGLQSYLLQFHGL VRLVHQERTLAFPLTIRCFLGCELPPEGSRAHVFFEVAVNGSSFVSFRPERALWQADTQVTSGVVTFTLQQLNAYNRTRYE LREFLEDTCVQYVQKHISAENTKGSQTSRSYTS

#### 靶点信息

研究背景	The protein encoded by this gene is a receptor for activated protein C, a serine protease activated by and involved in the blood coagulation pathway. The encoded protein is an N-glycosylated type I membrane protein that enhances the activation of protein C. Mutations in this gene have been associated with venous thromboembolism and myocardial infarction, as well as with late fetal loss during pregnancy. The encode d protein may also play a role in malarial infection and has been associated with cancer.
------	---

基因ID	10544
基因名	PROCR
Swiss	Q9UNN8
别名	PROCR;CCCA;CCD41;EPCR;h-PROCR

## 产品验证



Western blot analysis of PROCR expressed in Mouse kidney,MCF7,HepG2 using PROCR Rabbit pAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn