

NODAL Rabbit pAb

货号: **B14916**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IHC: Human glioma tissue
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	39kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,HepG2,293T,Rat liver
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 198-347 of human NOD AL (NP_060525.3).
序列	SNLSQEQRQLGGSTLLWEAESSWRAQEGQLSWEWGKRHRRHHLPDRSQLCRKVKFQVDFNLIGWGSWIIYPKQYNAY RCEGECPNPVGEEFHPTNHAYIQSLLKRYQPHRVPSTCCAPVKTKPLSMLYVDNGRVLLDHHKDMIVEECGCL

靶点信息

研究背景	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of pro teins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMA D family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate the mature protein, which regulates early embryonic development. This protein is required for maintenance of human embryonic stem cell pluripotency and may play a role in human place ntal development. Mutations in this gene are associated with heterotaxy, a condition characterized by ran dom orientation of visceral organs with respect to the left-right axis.
------	---

基因ID	4838
基因名	NODAL
Swiss	Q96S42
别名	NODAL;HTX5

产品验证



Western blot analysis of NODAL expressed in Rat liver,SH-SY5Y,HepG2 using NODAL Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn