

SMYD5 Rabbit pAb

货号: **B14931**

产品信息

| 反应 | Human,Mouse |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: HeLa |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 47kDa |
| 实测分子量 | 48kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HepG2,MCF7,HL-60,HeLa,SW480,Mouse skeletal muscle |
| 细胞定位 | |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant protein of human SMYD5 |
|------|------------------------------------|
| 序列 | |

靶点信息

| 研究背景 | Histone methyltransferase that specifically trimethylates 'Lys-20' of histone H4 to form trimethylated hist one H4 lysine 20 (H4K20me3 which represents a specific tag for epigenetic transcriptional repression (By similarity. In association with the NCoR corepressor complex, is involved in the repression of toll-like rece ptor 4 (TLR4-target inflammatory genes in macrophages by catalyzing the formation of H4K20me3 at the gene promoters (By similarity. Plays an important role in embryonic stem (ES cell self-renewal and differentiation (By similarity. Promotes ES cell maintenance by silencing differentiation genes through deposition of H4K20me3 marks (By similarity. Maintains genome stability of ES cells during differentiation through regulation of heterochromatin formation and repression of endogenous repetitive DNA elements by depositing H4K20me3 marks. |
|--------------|--|
| 基因 ID | 10322 |
| 基因名 | SMYD5 |
| Swiss | Q6GMV2 |
| 别名 | SMYD5;NN8-4AG;RAI15;RRG1;ZMYND23 |

产品验证



Western blot analysis of SMYD5 expressed in Mouse skeletal muscle, HepG2, MCF7 using SMYD5 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug pe r lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn